


What Is Erectile Dysfunction (ED)?

ED is defined as the persistent inability to achieve or maintain an erection that is firm enough to have sexual intercourse.¹

How Common Is ED?

ED is a common problem and it's important to know you're not alone.

Age breakdown of men with total inability to achieve an erection¹



To learn more about ED visit **EDTreatments.com**



Answers for Men

What Causes ED?^{1,3}

There's no single cause of ED. There are real physical and psychological reasons for ED.

Some common causes are:

- Diabetes
- Cardiovascular problems (high blood pressure, heart disease)
- Prostate cancer treatment
- Surgery (prostate, bladder, colon, rectal)
- Medications (blood pressure, antidepressants)
- Lifestyle choices (smoking, excessive alcohol, obesity, lack of exercise)
- Spinal cord injuries
- Hormone problems

ED is not just a medical issue; it can also deeply affect relationships.⁴



How did ED affect your relationship?

"The intimacy that we used to have went away. All of a sudden, it was like we were completely separated. There was no connection."

-Tom

Erections and How They Work³

To learn more about ED, it is important to understand how the penis normally works. The erection process includes the following 5 stages:

1. Initial Filling

With psychological or sexual stimulation, neurotransmitters cause penile smooth muscles to relax, increasing blood flow to the corporal bodies (the 2 chambers within the shaft of the penis).



2. Partial Erection


Penile arteries expand to accommodate the increased blood flow needed to elongate and expand the penis.






3. Full Erection

The increased volume of blood within the penis is prevented from draining, thus expanding the penis to full erection.



4. Rigid Erection

Maximum rigidity is attained. The tip of the penis and penis body enlarge until penile veins are forcefully compressed. This increases engorgement and maintains maximum penile rigidity.



5. Return to Flaccidity

Muscle contractions result in increased blood outflow from the penis, thus decreasing penile length and girth until flaccid.



ED Treatment Options³

The good news is that there are treatment options for ED, and there are multiple options available. For some men, oral medications don't work, so it's important to know all of your options. ⁵⁻⁷ Your doctor will discuss options that may be appropriate for you, along with the potential risks and benefits.

Oral Medications

There are a number of prescription medications available that may improve blood flow to the penis. Combined with sexual stimulation, this can produce an erection.



Vacuum Erection Devices

A plastic cylinder is placed over the penis, and a pump (either manual or battery operated) creates a vacuum suction within the cylinder, drawing blood into the penis to create an erection.


A stretchable tension band is placed at the base of the penis to help maintain the erection.





Intraurethral Suppositories

An applicator containing a small pellet (suppository) is inserted into the urethra and the pellet is released. The pellet dissolves and increases blood flow to the penis, creating an erection.



Injections

With injection therapy, a needle is used to inject medication directly into the penis. The medication allows blood to flow into the penis, creating an erection



Penile Implants

A medical device is implanted into the penis that allows men with ED to achieve an erection. This device is entirely contained within the body. To operate, one squeezes and releases the pump in the scrotum to achieve an erection.

What is a Penile Implant?

A penile implant is a medical device that is implanted into a man's body. The surgery is typically performed as an inpatient procedure.¹⁴

Penile implants have been in clinical use for over 45 years⁸ and Boston Scientific devices have helped over 500,000 men return to an active and satisfying sex life.⁹ Implants provide an option for men who have tried other treatments without success. Many studies show most patients and their partners are highly satisfied with the results.¹⁰

Finding a treatment option for ED can be a life-changing event for many men and their partners. Choosing the penile implant that is best for you is a very personal decision. Because each type of implant offers unique features, you will want to discuss the risks and benefits of each with your doctor to help you choose the option that is right for you.





Features of a Penile Implant:

- Entirely contained in the body¹⁰
- May offer a long-term treatment option to ED¹⁵
- Once activated, you can maintain an erection as long as you desire¹⁵
- Spontaneous have sex when the mood strikes¹⁵
- Typically does not interfere with ejaculation or orgasm¹⁵
- Designed to feel natural during intercourse¹⁵

Risks of a Penile Implant¹⁰:

- Will make natural or spontaneous erections as well as other interventional treatment options impossible
- If an infection occurs, the implant may have to be removed
- May cause the penis to become shorter, curved, or scarred
- Pain (typically associated with the healing process)
- There may be mechanical failure of the implant


Patient and Partner Satisfaction

ED can limit your intimacy, affect your self-esteem, and impact your most important relationships.⁴

Many studies show that penile implants may offer the satisfying results that so many couples seek. Consult your doctor to determine which type of implant is best suited for your condition and lifestyle.

In one study of 106 men with erectile dysfunction, 32 received a penile implant. Of these patients, 93% were satisfied with their implant.¹¹

Overall Patient Satisfaction with erectile dysfunction treatments¹¹




To hear real patient stories visit **EDTreatments.com**




Often patients wonder if their sexual activity will be the same and if they will be able to experience orgasms and sensation. In one study of 200


patients and 120 partners, both men and their partners found the AMS 700™ penile implant to be satisfying¹2:



92% of patients reported sexual activity with the implant to be excellent or satisfactory



96% of their partners reported sexual activity with the implant to be excellent or satisfactory



75% of patients reported no change in their orgasm, and **20% reported** better orgasms, following the surgery



50% of patients reported their erections as satisfactory, and48% reported excellent erections, following the surgery

To hear real patient stories visit **EDTreatments.com**

AMS 700™ Penile Prosthesis – Three-Piece Inflatable 1

The 3-piece, fluid-filled, inflatable penile implant consists of a pair of cylinders that are implanted into the penis, a pump that is implanted in the scrotum, and a reservoir that is implanted into the lower abdomen.

Squeezing and releasing the pump moves fluid into the cylinders, creating an erection. Deflate the device by pressing the deflate button on the pump. The penis then returns to a soft, flaccid and natural-looking state.¹⁶

The 3-piece penile implant is entirely contained in the body and is designed to produce a natural erection.

Another feature of the 3-piece penile implant is that it expands in girth (all AMS 700™ cylinders) and length (AMS 700 LGX™ cylinders).

Additional Implant Options

Boston Scientific offers 2 additional types of penile implants, the Ambicor™ 2-Piece Inflatable Penile Prosthesis and the Tactra™ 1-Piece Malleable (Bendable) Penile Prosthesis.

Please talk to your doctor to determine which implant is right for you.

e Penile Implant



AMS 700[™] LGX Penile Implant

- The only penile implant on the market with cylinders designed to expand in girth and length of 10-25%¹⁷
- Cylinders designed to address the #1 concern men with ED have – loss of penile length^{18,19}
- Controlled expansion is designed to provide maximum rigidity and optimal length and girth expansion¹⁷
- Firm, rigid erection that can last as long as desired¹⁷
- Natural, flaccid appearance when deflated¹⁷


AMS 700[™] CX Penile Implant

- Controlled expansion is designed to provide maximum rigidity and optimal girth expansion²⁰
- Firm, rigid erection that can last as long as desired¹⁷
- Natural, flaccid appearance when deflated¹⁷

AMS 700™ CXR Penile Implant

- All the benefits of the AMS 700 implant series, including girth expansion for patients who may require shorter and narrower cylinders²⁰
- Controlled expansion is designed to provide maximum rigidity and optimal girth expansion¹⁷
- Firm, rigid erection that can last as long as desired¹⁷
- Natural, flaccid appearance when deflated¹⁷

AMS 700[™] Penile Prosthesis



With So Much to Know, What Haven't I Thought of?

Surgery

The implant procedure is typically done on an inpatient basis. Patients generally check in the morning of the procedure and are discharged within 24 hours. Patients receive anesthesia during the surgery. Your doctor can tell you more about the surgical procedure.

Healing

You may experience pain and soreness at the surgical site. Typically, it takes a few days to return to your regular daily routine of light activity. The full healing time before using your device is usually about 6 weeks.



Sexual Health Inventory for Men (SHIM)³

This quiz is often used to help doctors determine a patient's degree of erectile dysfunction (ED).

Discuss your answers with your doctor today.

After you take the quiz make a list of your questions and concerns, clip out your completed quiz, and return to the urologist's office for a more in-depth conversation about possible next steps.

Over the Past 6 Months

How do you rate your confidence that you		Very Low	Low	Moderate	High	Very High
could get and keep an erection?	0	1	2	3	4	5
2. When you had erections with sexual stimulation, how often were your erections hard enough for penetration (entering your partner)?	No Sexual Activity O	Almost Never or Never	A Few Times (Much Less Than Half the Time)	Sometimes (About Half the Time)	Most Times (Much More Than Half the Time)	Almost Always or Always
3. During sexual intercourse, how often were you able to maintain your erection after you had penetrated (entered) your partner?	Did Not Attempt Intercourse	Almost Never or Never	A Few Times (Much Less Than Half the Time)	Sometimes (About Half the Time)	Most Times (Much More Than Half the Time)	Almost Always or Always
4. During sexual intercourse, how difficult was it to maintain your erection to completion of intercourse?	Did Not Attempt Intercourse	Extremely Difficult 1	Very Difficult 2	Difficult 3	Slightly Difficult 4	Not Difficult 5
5. When you attempted sexual intercourse, how often was it satisfactory for you?	Did Not Attempt Intercourse	Almost Never or Never	A Few Times (Much Less Than Half the Time)	Sometimes (About Half the Time)	Most Times (Much More Than Half the Time)	Almost Always or Always

Add the numbers corresponding to questions 1-5. Total:

The Sexual Health Inventory for Men further classifies ED severity with the following breakpoints:

1-7 Severe ED 8-11 Moderate ED 12-16 Mild to Moderate ED 17-21 Mild ED

Questions and Concerns



AMS 700™ Series Inflatable Penile Prosthesis Brief Summary

The AMS 700™ Series Inflatable Penile Prosthesis is intended for use in the treatment of chronic, organic, male erectile dysfunction (impotence).

These devices are contraindicated in patients who have active urogenital infections or active skin infections in the region of surgery or (for the AMS 700 with InhibiZone™) have a known sensitivity or allergy to rifampin, minocycline, or other tetracyclines.

Implantation will make latent natural or spontaneous erections, as well as other interventional treatment options, impossible. Men with diabetes, spinal cord injuries or open sores may have an increased risk of infection. Failure to evaluate and treat device erosion may result in infection and loss of tissue. Implantation may result in penile shortening, curvature, or scarring.

Possible adverse events include, but are not limited to, urogenital pain (usually associated with healing), urogenital edema, urogenital ecchymosis, urogenital erythema, reservoir encapsulation, patient dissatisfaction, auto-inflation, mechanical malfunction, and impaired urination.

Prior to using these devices, please review the Instructions for Use for a complete listing of indications, contraindications, warnings, precautions, and potential adverse events.

Ambicor™ Penile Prosthesis Brief Summary

The AMS Ambicor™ Penile Prosthesis is intended for use in the treatment of chronic, organic, male erectile dysfunction (impotence).

These devices are contraindicated in patients who have active urogenital infections or active skin infections in the region of surgery.

Implantation will make latent natural or spontaneous erections, as well as other interventional treatment options, impossible. Men with diabetes, spinal cord injuries, or open sores may have an increased risk of infection. Failure to evaluate and treat device erosion may result in infection and loss of tissue. Implantation may result in penile shortening, curvature, or scarring.

Possible adverse events include, but are not limited to, urogenital pain (usually associated with healing), patient dissatisfaction, mechanical malfunction, auto-inflation, penile curvature or sensation change, urogenital hematoma, urogenital edema, and infection.

Prior to using these devices, please review the Instructions for Use for a complete listing of indications, contraindications, warnings, precautions, and potential adverse events.



Patient Testimonials

The stories throughout this brochure recount the experiences of people who are using AMS therapies related to erectile dysfunction. AMS invited these people to share their stories candidly. As you review them, please bear in mind that the experiences are specific to these particular people. As with all medical treatment, not every response is the same – results vary.

Take the next step

Erectile dysfunction (ED) can limit your intimacy, affect your self esteem and have an impact on your relationship with your partner. But there is good news – today, nearly every case of ED is treatable.

To learn more about treatments that are available visit **EDTreatments.com**



Talk with a urologist who specialises in ED and offers the full range of treatment options, including implants.



Need help finding an experienced doctor near you?

Visit our "Find an Implanter" feature at EDTreatments.com



Watch real patient stories at EDTreatments.com

Notes

References

- Erectile dysfunction. http://kidney.niddk.nih.gov/ KUDiseases/pubs/ED/ErectileDysfunction_508.pdf. National Kidney and Urologic Diseases Information Clearinghouse Web site. Downloaded January 21, 2013.
- Selvin E, Burnett A, Platz E. Prevalence and risk factors for erectile dysfunction in the U.S. Am J Med. 2007:(120):151-157
- Erectile dysfunction (ED). www.auanet.org/content/ education-and-meetings/med-stu-curriculum/ed.pdf. American Urological Association Web site. Downloaded January 21, 2013.
- DiMeo PJ. Psychosocial and relationship issues in men with erectile dysfunction. Urol Nurs. 2006;26(6):442-453.
- 5. Viagra® Prescribing Information, Pfizer Inc. Revised January 2010.
- Cialis* Prescribing Information, Lilly USA, LLC. Revised October 2011.
- Levitra® Prescribing Information, Bayer HealthCare Pharmaceuticals. Revised November 2011.
- 8. Scott FB, Bradley WE, Timm GW. Management of erectile impotence: use of implantable inflatable prosthesis. Urology. 1973;2(1):80-82.
- Penile prosthesis. http://www.essm.org/society/ esha/malesd/oraltreatments/penileprosthesis.html. European Society for Sexual Medicine (ESSM) Web site. Downloaded January 1, 2013.
- AMS 700™ Patient Manual. Information and Instructions for Patients Considering an Inflatable Penile Prostheses. American Medical Systems. 2012.

- 11. Rajpurkar A, Dhabuwala C. Comparison of satisfaction rates and erectile function. J Urol. 2003:170:159-163.
- Montorsi F, Rigatti P, Carmignani G, et al. AMS three-piece inflatable implants for erectile dysfunction: a long-term multi-institution study in 200 consecutive patients. Eur Urol. 2000:37:50-55.
- 13. Dymedex Consulting. Assessment of the market potential for Penile Prosthetic Implants (PPI). November, 2010.
- AMS 700™ Penile Prosthesis Operating Room Manual. American Medical Systems. 2012.
- 15. AMS 700™ Penile Prosthesis Product Line Instructions for Use. American Medical Systems. 2012.
- Otero JR, Cruz CR, Gómez BG, et al. Comparison of the patient and partner satisfaction with 700CX and Titan penile prostheses. Asian J Androl. 2017 May-Jun;19(3):321-5.
- 17. Data on file with Boston Scientific and based on market research by Dymedex.
- Sellers T, Dineen M, Salem E, et al. Vacuum preparation, optimization of cylinder length and postoperative daily inflation reduces complaints of shortened penile length following implantation of inflatable penile prosthesis. Adv in Sex Med. 2013 Jan;3(91):14-8.
- Kamel I, Gadalla A, Ghanem H, et al. Comparing penile measurements in normal and erectile dysfunction subjects. J Sex Med. 2009 Aug;6(8):2305-10.
- 20. AMS 700[™] with MS Pump[™] Penile Prosthesis Operating Room Manual. American Medical Systems. 2018.



All cited trademarks are the property of their respective owners.

CAUTION: Law restricts these devices to sale by or on the order of a physician. Indications, contraindications, warnings and instructions for use can be found in the product labelling supplied with each device. Information for the use only in countries with applicable health authority product registrations.

UROPH-413712-AB DEC 2020. Produced by Gosling.

www.bostonscientific.eu

© 2021 Boston Scientific Corporation or its affiliates. All rights reserved. DINURO2230EC